

1997

## Dedication: Arnel R. Hallauer, Scientist, Maize Breeder, Quantitative Geneticist

Kendall R. Lamkey

*U.S. Department of Agriculture, [krlamkey@iastate.edu](mailto:krlamkey@iastate.edu)*

Follow this and additional works at: [http://lib.dr.iastate.edu/agron\\_pubs](http://lib.dr.iastate.edu/agron_pubs)

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Plant Breeding and Genetics Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/agron\\_pubs/260](http://lib.dr.iastate.edu/agron_pubs/260). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Book Chapter is brought to you for free and open access by the Agronomy at Iowa State University Digital Repository. It has been accepted for inclusion in Agronomy Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Dedication: Arnel R. Hallauer, Scientist, Maize Breeder, Quantitative Geneticist

## Abstract

Arnel R. Hallauer has dedicated his life to breeding; therefore, it is appropriate that this issue of Plant Breeding Reviews be dedicated to him. Dr. Hallauer's primary contributions have been in understanding quantitative inheritance in maize, developing breeding methodology, evaluating and utilizing recurrent selection for population improvement, and developing inbred lines for use in maize hybrids. Dr. Hallauer has also been very involved in graduate student education. The success of his students in academia and industry is a measure of his ability as a graduate educator and represents one of his greatest contributions to plant breeding. He has also been extensively involved in international activities and is known throughout the world for the impact of his efforts in that arena. Sprague and Lamkey (1992) have reviewed Dr. Hallauer's career in the context of the development of quantitative genetics and his research career will be emphasized here.

## Disciplines

Agricultural Science | Agronomy and Crop Sciences | Plant Breeding and Genetics

## Comments

This is a chapter from Lamkey, K. R. (1997) Dedication: Arnel R. Hallauer, Scientist, Maize Breeder, Quantitative Geneticist, in Plant Breeding Reviews, Volume 15 (ed J. Janick), John Wiley & Sons, Inc., Oxford, UK. doi: [10.1002/9780470650097.ch1](https://doi.org/10.1002/9780470650097.ch1). Posted with permission.

## Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

## Dedication:

Arnel R. Hallauer  
Scientist, Maize Breeder,  
Quantitative Geneticist\*

*Kendall R. Lamkey*

Field Crops Research Unit  
United States Department of Agriculture  
Agricultural Research Service  
Department of Agronomy  
Iowa State University  
Ames, Iowa 50011

Arnel R. Hallauer has dedicated his life to breeding; therefore, it is appropriate that this issue of *Plant Breeding Reviews* be dedicated to him. Dr. Hallauer's primary contributions have been in understanding quantitative inheritance in maize, developing breeding methodology, evaluating and utilizing recurrent selection for population improvement, and developing inbred lines for use in maize hybrids. Dr. Hallauer has also been very involved in graduate student education. The success of his students in academia and industry is a measure of his ability as a graduate educator and represents one of his greatest contributions to plant breeding. He has also been extensively involved in international activities and is known throughout the world for the impact of his efforts in that arena. Sprague and Lamkey (1992) have reviewed Dr. Hallauer's career in the

\*Joint contribution from the Field Crops Research Unit, USDA, Agricultural Research Service, Dept. of Agronomy, Iowa State Univ. and Journal Paper No. J-17065 of the Iowa Agric. and Home Economics Exp. Sta. Project No. 3082.

---

*Plant Breeding Reviews, Volume 15* Edited by Jules Janick

© 1997 John Wiley & Sons, Inc. ISBN: 978-0-471-18904-6

context of the development of quantitative genetics and his research career will be emphasized here.

## BIOGRAPHICAL SKETCH

Arnel Roy Hallauer was born on May 4, 1932, at Netawaka, Kansas. His interest in maize research began in the fall of 1946, when at age 14 he had a part-time job with Dr. Lloyd A. Tatum harvesting experimental maize trials in northeast Kansas. He continued working part time with Dr. Tatum while attending high school (1946–1950), planting experimental maize trials in the spring and harvesting them in the fall. During the summer months of his high school years, he detasseled maize for the production of double-cross hybrid seed for Mr. Carl Overly of the Kansas Crop Improvement Association. These associations with Dr. Tatum and Mr. Overly led him to Kansas State University where he received his B.S. degree with honors in 1954 with a major in plant science. During his undergraduate studies, he was continuously employed part time with the cooperative federal-state maize-breeding project directed by Dr. Tatum at Kansas State University. There he experienced all aspects of the breeding project including preparing seed, planting, collecting data, harvesting, pollinating in the breeding nurseries, and analyzing and summarizing data of experimental trials.

After finishing his undergraduate degree, Arnel served for two years in the military (1954–1956). Dr. Tatum, impressed with his young assistant's dedication and work ethic, urged Arnel to attend graduate school for an advanced degree in plant breeding. After considering offers from many universities, he chose Iowa State because the offer included the opportunity to work with Dr. George F. Sprague, a maize breeder and geneticist who had a highly-regarded research program. Dr. Hallauer completed the requirements for the M.S. degree in plant breeding in 1958. While working for the USDA-ARS, he continued in graduate school and received his Ph.D. in 1960. Dr. W. A. Russell served as his major advisor for the Ph.D. degree after Dr. Sprague was transferred to Beltsville, Maryland, in 1958 to become the USDA-ARS Investigations Leader for maize and sorghum research.

With the completion of his graduate training, Dr. Hallauer was transferred to North Carolina State College as a USDA-ARS postdoctoral research geneticist to spend a year with Professors H. C. Robinson and R. H. Moll. In 1962, he was transferred back to Ames as a USDA-ARS research geneticist; he continued in this capacity until 1989. In December of 1989, Dr. Hallauer retired from the USDA-ARS and accepted a posi-

tion with Iowa State University as professor of plant breeding. In 1991, he was named the C. F. Curtiss Distinguished Professor in Agriculture.

## QUANTITATIVE GENETICS

Dr. Hallauer has devoted his career to understanding the inheritance of agronomic traits in maize. Because most traits of importance in maize are controlled by many genes, Dr. Hallauer was mainly interested in the quantitative genetics of maize populations. The main questions of interest were the types of gene action controlling quantitative traits, the amounts and types of genetic variance in maize populations (additive, dominance, or epistatic), and the heritability of agronomic traits. These questions were of utmost importance because they were directly related to the rate of genetic gain that could be expected from recurrent selection in maize populations.

Dr. Hallauer began his career studying the inheritance of grain moisture at harvest and time of flowering in maize. He then focused his attention on estimating the genetic variance components for agronomic traits in open-pollinated populations and synthetic cultivars of maize. He found that additive genetic variance frequently accounted for the majority of the total genetic variation for grain yield, and dominance variance, although much smaller, was usually significant and accounted for the remainder. Extensive studies to detect epistatic genetic variance in maize synthetics revealed that it was a minor contributor to the total genetic variance for grain yield.

The one major exception to the pattern of genetic variances observed by Dr. Hallauer was Iowa Stiff Stalk Synthetic (BSSS). Variance component studies in BSSS have consistently shown that the additive and dominance variance components are equal in magnitude. This result was in direct contrast to average results from other maize populations where the additive variance was usually twice the dominance variance (Hallauer and Miranda 1988). This result must be directly related to the choice of inbred lines used to form BSSS. It is also interesting to note that BSSS is one of the most important sources of inbred lines in the maize industry.

One of the most striking phenotypic observations in maize is the extensive inbreeding depression that is observed for grain yield when maize is self-pollinated. The studies that Dr. Hallauer conducted on inbreeding depression in maize were among the first and are now classics (Hallauer and Sears 1975; Good and Hallauer 1977). These studies demonstrated that inbreeding depression for grain yield was a linear function of the

inbreeding coefficient. Linear regression usually accounted for 93 to 99% of the variation among means of the inbred generations. The linearity of inbreeding depression was additional evidence in support of conclusions from the variance component studies that epistasis was not important in maize synthetics.

Because of the importance of directional dominance in maize as evidenced by the inbreeding depression observed for grain yield, there has been extensive research into tester theory and the relationship between line and hybrid performance. The choice of testers in a maize improvement program is one of the more important decisions a breeder makes in a breeding program. The choice of a tester is highly dependent on the level of dominance (partial to complete dominance vs. overdominance) and the frequency of favorable alleles in the tester population (or line) in comparison to the population being tested. The theory shows that testers with a low frequency of favorable alleles will maximize genetic variance among testcross progenies for all levels of dominance. The problem is that maize breeders are reluctant to use low-yielding inbreds as testers and inbreds that are low yielding may not necessarily have low frequencies of alleles for grain yield. Hallauer and Lopez-Perez (1979) conducted an extensive study evaluating the discriminatory power of five types of testers selected for their expected differences in allelic frequency for grain yield. The five testers were crossed to 50  $S_1$  lines and the 50  $S_8$  lines derived from these  $S_1$  lines. Four of the testers were related to the population the lines were derived from and one of the testers was unrelated. Hallauer and Lopez-Perez found that genetic variance among testcrosses ranked according to the expected frequency of the favorable allele for grain yield. Genetic variances among testcrosses to the unrelated tester were similar to those found with the lowest yielding (and lowest expected frequency of the favorable allele) tester. These results were a significant finding because they demonstrated that unrelated testers (usually from the opposite heterotic group) have discriminatory power similar to using a low-yielding tester.

Quantitative genetics has undoubtedly contributed substantially to maize breeding methodology. Dr. Hallauer and J. B. Miranda, Filho have done an excellent job of summarizing quantitative genetics in maize in their book entitled *Quantitative Genetics and Maize Breeding*. The book is an extensive summary of quantitative genetic studies in maize and relates this information to the various breeding methods used in maize. The book is in the second edition and is in high demand by breeders throughout the world.

## BREEDING METHODOLOGY

Dr. Hallauer's contributions to breeding methodology were influenced in part by the debate in the 1940s and 1950s over the importance of dominance and overdominance in the expression of heterosis for grain yield. Comstock et al. (1949) proposed half-sib reciprocal recurrent selection in response to this debate because this breeding method took advantage of all types of gene effects (including epistasis) and was only slightly less efficient than other methods when overdominance and/or epistasis were not important. Cockerham (1961) evaluated the expected genetic variance among unrelated single, three-way, and double crosses and demonstrated that the genetic variance among single crosses was always greater than among three-way crosses and that the genetic variance among three-way crosses was always greater than among double crosses. If only additive effects are important, then selection among single crosses would be twice as effective as selection among double crosses, and if nonadditive effects (dominance and epistasis) are important, selection among single crosses would be four times as effective as selection among double crosses. These theoretical developments and other concurrent developments paved the way for Dr. Hallauer's ideas on breeding methodology for developing superior maize single crosses.

Hallauer (1967a; 1967b) proposed a procedure for developing single-cross hybrids that maximized selection for nonadditive effects. The procedure required the use of two-eared maize populations. In the first phase, crosses are made between individual  $S_0$  plants from two populations. At the same time the crosses are made, the  $S_0$  plants are also selfed. In this way, both hybrids and  $S_1$  lines could be developed from the same plants. The following year, the  $S_0 \times S_0$  hybrids are evaluated in yield trials and the pairs of  $S_1$  lines are grown in the nursery for further crossing and selfing. Dr. Hallauer recommended that four to six crosses be produced within each pair of  $S_1$  lines. The procedure is then repeated until pure-line hybrids are developed. At each stage of evaluation, only the best hybrids are retained. In a comparison of  $S_0 \times S_0$  crosses with the  $S_1 \times S_1$  crosses, Dr. Hallauer showed that the first generation of selection was effective for improving grain yield. Hallauer (1967b) also outlined a procedure where one-eared source populations could be used.

Hallauer and Eberhart (1970) suggested a modification of the two-eared method for developing single crosses for population improvement. They called this method reciprocal full-sib selection to distinguish it from the reciprocal recurrent selection proposed by Comstock et al.

(1949). The method was to cross pairs of  $S_0$  plants (one plant from each of two populations) and simultaneously self the plants. The  $S_0 \times S_0$  hybrids are evaluated in yield trials, the best are selected, and remnant  $S_1$  seed is used to intermate to form the two improved populations for continued selection. This breeding procedure incorporated yield testing of  $S_0 \times S_0$  hybrids for population improvement as well as early testing of single-cross combinations. Thus, the best selections are used to form a new population for further selection and they are incorporated directly into a single-cross development program. One of the main advantages over reciprocal recurrent selection is that twice as many plants from the source populations can be evaluated using the same amount of resources required in reciprocal recurrent selection.

## GERMPLASM DEVELOPMENT AND IMPROVEMENT

Much of Dr. Hallauer's efforts have been devoted to germplasm development and population improvement via recurrent selection. Dr. Hallauer works with both adapted temperate and exotic germplasm. He currently has 10 active recurrent selection programs involving 12 populations. The primary traits under selection are grain yield, grain moisture, root lodging, and stalk lodging. Other traits under selection in specific programs are ear length, maturity, and resistance to European corn borer (*Ostrinia nubilalis* Hübner). Once populations reach a desired level of performance, the improved populations are released to the general public and registered in *Crop Science*.

The progenies selected to form each new cycle of a population are routinely placed in a pedigree breeding program to assess their potential as parents of hybrids. This feature has been a hallmark of the Cooperative Federal-State Breeding Program that has resulted in the release of many influential inbred lines. Currently and traditionally, the responsibility for managing the pedigree breeding program has been with the state breeder. Dr. Hallauer currently has this responsibility, replacing Dr. W. A. Russell who retired in 1989. Inbred lines resulting from this program are released as parental lines to be used in public and private breeding programs. Although some may view this as competition with the private sector, that is not the intent and is one of the reasons that hybrid recommendations are never made. The development and release of parental lines is a natural extension of recurrent selection and is a way of transferring improved germplasm to the general public. The pedigree breeding program is also an invaluable aid in educating and training graduate students and monitoring progress in recurrent selection programs. It is



noteworthy to point out that commercial germplasm has never been incorporated into any of the populations or inbred lines that have been released.

Dr. Hallauer has had great success in using recurrent selection to incorporate exotic germplasm into his breeding program. In his first work with tropical germplasm, Dr. Hallauer compared two methods of adapting "Eto Composite," a tropical maize population from Colombia, South America, to temperate climates. In the first method, he used mass selection for earlier silking and adaptation. In the second method, he crossed "Eto Composite" with six early inbred lines and intermated the resulting  $F_1$ 's to form a new synthetic. He demonstrated that mass selection for adaptation and earlier silking was an effective method for adapting exotic germplasm to temperate environments. In the "Eto Composite," time to silking was decreased an average of four days per year with mass selection and required only five years to adapt "Eto Composite" to Iowa. This was an important finding because it allows the utilization of populations with 100% exotic germplasm, which increases the probability that novel genotypes will be derived from the population. The second method Dr. Hallauer used to adapt "Eto Composite" to Iowa involved the incorporation of corn-belt germplasm into the exotic germplasm, which raises issues about genetic recombination between the adapted and exotic genomes. Dr. Hallauer's research demonstrated that corn-belt germplasm does not need to be incorporated into tropical germplasm in order to adapt it to temperate environments.

By using the mass-selection technique, Dr. Hallauer has successfully adapted "Eto Composite," "Tuxpeño" (tropical germplasm obtained from CIMMYT in Mexico), "Antigua" (tropical germplasm obtained from CIMMYT in Mexico), and "Suwan-1" (tropical maize population developed in Thailand). Recurrent selection is ongoing in all of these populations except for "Antigua." Dr. Hallauer is currently adapting "Tusón" (tropical germplasm from the Caribbean and South America) and is converting adapted versions of "Mexican Dent" and "Cateto" (tropical germplasm from South America)  $\times$  "Caribbean Flint" from white to yellow kernels. These populations have all been used extensively in their native environments. The adapted versions of these populations exhibit excellent yield potential and a greater incidence of disease and insect resistance than most corn-belt populations.

Dr. Hallauer has been involved with and participated in the release of many populations, synthetics, and inbred lines. He has been unselfish with his germplasm and is committed to the free exchange of germplasm. The dedication and commitment of Dr. Hallauer and other members of the Cooperative Federal-State Maize Breeding project in Ames to the free

exchange of germplasm has made Ames one of the top centers in the world for the development and dissemination of maize germplasm.

## GRADUATE EDUCATION

Dr. Hallauer is dedicated to graduate education. In 1966, he created a special topics course on quantitative genetics, which in 1969 was incorporated into the Department of Statistics where it has been taught ever since. He conducted a spring semester plant breeding seminar from 1975 to 1993, has taught advanced plant breeding since 1986, and taught field plot techniques in plant breeding from 1989 to 1994.

Dr. Hallauer has devoted much of his time to the training of graduate students. His students have come from the United States and 18 foreign countries. He has been major advisor to 75 students (51 Ph.D. and 24 M.S.) and has hosted 25 visiting scientists or postdoctoral students from 11 different countries. At Iowa State, he has served on more than 100 graduate student program of study committees. A host of domestic and foreign visitors have visited Dr. Hallauer and the maize breeding project.

## HONORS AND AWARDS

Dr. Hallauer has been extensively recognized for his achievements and contributions to plant breeding and science. Within the state of Iowa, he received the Applied Research and Extension Award (ISU, 1981), the Distinguished Fellow Award (Iowa Academy of Sciences, 1995), a Faculty Citation Award (Iowa State Alumni Association, 1987), the Distinguished Service to Agriculture Award (Gamma Sigma Delta, ISU, 1990), a Governor's Science Award (Terry Branstad, Governor of Iowa, 1990), the C. F. Curtiss Distinguished Professor in Agriculture Award (ISU, 1991), the Burlington Northern Career Achievement in Research Award (ISU Foundation, 1991), the Henry A. Wallace Award for Distinguished Service to Agriculture (ISU Alumni Association, 1992), and a Distinguished Achievement Citation (ISU Alumni Association, 1995); he was elected an honorary member of the Iowa Crop Improvement Association (1995).

At the national level, Dr. Hallauer has been a member of the American Society of Agronomy (ASA) since 1958 and the Crops Science Society of America (CSSA) since 1961. He has served these societies in many ways and was elected a fellow of ASA (1979) and a fellow of CSSA (1985). He received the Crop Science Award (CSSA, 1981), the Agro-

nomic Achievement Award—Crops (ASA, 1989), the Dekalb Crop Science Distinguished Career Award (CSSA, 1990), and the Agronomic Research Award (ASA, 1992). The commercial sector has recognized his achievements by awarding him the Northrup-King Recognition Award for Research in Corn Breeding (1984), the National Council of Commercial Plant Breeder's Genetics and Breeding Award (1984), and the National Agri-Marketing Association's Excellence in Research Award (1993). For his outstanding service to the USDA-ARS, Dr. Hallauer received the Scientist of the Year Award (1985) and was elected to the USDA-ARS Science Hall of Fame (1992). His crowning achievement, and one that has been attained by only a few maize breeders, was his election to the U.S. National Academy of Sciences.

## THE MAN

Dr. Hallauer is a self-effacing man. Despite his achievements and recognition, he has remained humble and modest. He treats everyone as an equal and is always willing to listen, share his knowledge and germplasm, and help out in what ever way he can. He is a prolific writer, an outstanding editor, an avid reader, and an excellent speaker. For enjoyment, he reads books on history, including science history and U.S. and world history, and popular fiction. He has a great sense of humor and has that rare ability to incorporate humor into his seminars. He is a people person and makes friends easily. Arnel Hallauer is a man who has the distinction of being respected, admired, and loved by all.

## LITERATURE CITED

- Cockerham, C. C. 1961. Implication of genetic variances in a hybrid breeding program. *Crop Sci.* 1:47–52.
- Comstock, R. E., H. F. Robinson, and P. H. Harvey. 1949. A breeding procedure designed to make maximum use of both general and specific combining ability. *Agron. J.* 41:360–367.
- Good, R. L., and A. R. Hallauer. 1977. Inbreeding depression in maize by selfing and full-sibbing. *Crop Sci.* 17:935–940.
- Hallauer, A. R. 1967a. Development of single-cross hybrids from two-eared maize populations. *Crop Sci.* 7:192–195.
- Hallauer, A. R. 1967b. Performance of single crossed hybrids developed from two-eared varieties. p. 74–81. In: *Proc. 22nd Annu. Hybrid Corn Industry Research Conference*, Chicago, IL, 13–14 Dec. 1967. Am. Seed Trade Assoc., Washington, D.C.
- Hallauer, A. R., and E. Lopez-perez. 1979. Comparisons among tasters for evaluating lines of corn. p. 57–75. In: *Proc. 34th Annu. Corn Sorghum Research Conference*, Chicago, IL. Am. Seed Trade Assoc., Washington, D.C.
- Hallauer, A. R., and J. B. Miranda, Fo. 1988. *Quantitative genetics and maize breeding*, 2nd edition. Iowa State Univ. Press, Ames.

- Hallauer, A. R., and J. H. Sears. 1972. Integrating exotic germplasm into Corn Belt maize breeding programs. *Crop Sci.* 12:203–206.
- Sprague, G. F., and K. R. Lamkey. 1992. Arnel R. Hallauer: An appreciation. *Maydica* 37:1–6.

## SELECTED PUBLICATIONS OF A. R. HALLAUER

### Research Papers

- Hallauer, A. R., and W. A. Russell. 1961. Effects of some selected weather factors on grain moisture reduction from silking to physiological maturity in corn. *Agron. J.* 53:225–229.
- Hallauer, A. R., and W. A. Russell. 1962. Estimates of maturity and its inheritance in maize. *Crop Sci.* 2:289–294.
- Scott, G. E., A. R. Hallauer, and F. F. Dicke. 1964. Types of action conditioning resistance to European corn borer leaf feeding. *Crop Sci.* 4:604–606.
- Hallauer, A. R. 1964. Estimation of soil variability and convenient plot size from corn trials. *Agron. J.* 56:493–499.
- Hallauer, A. R. 1965. Inheritance of flowering in maize. *Genetics* 52:129–137.
- Hallauer, A. R., and J. H. Sears. 1966. Influence of time of day and silk treatment on seed set in maize. *Crop Sci.* 6:216–218.
- Schmidt, J. L., and A. R. Hallauer. 1966. Estimating harvest date of corn in the field. *Crop Sci.* 6:227–331.
- Hallauer, A. R., and S. A. Eberhart. 1966. Evaluation of synthetic varieties of maize for yield. *Crop Sci.* 6:423–427.
- Hallauer, A. R. 1967. Development of single-cross hybrids from two-eared maize populations. *Crop Sci.* 7:192–195.
- Hallauer, A. R. 1967. Performance of single-cross hybrids developed from two-ear varieties. *Hybrid Corn Industry Res. Conf. Proc.* 22:74–81.
- Hallauer, A. R., and J. A. Wright. 1967. Genetic variances in the openpollinated variety of maize, Iowa Ideal. *Zuchter* 37:178–185.
- Hallauer, A. R., C. D. Hutchcroft, M. T. Hillson, and R. L. Higgs. 1967. Relation among three maturity measurements and yield of grain in corn. *Iowa State J. Sci.* 42:121–136.
- Eberhart, S. A., and A. R. Hallauer. 1968. Genetic effects for yield in single, three-way, and double-cross maize hybrids. *Crop Sci.* 8:377–379.
- Hallauer, A. R., and J. H. Sears. 1968. Second phase in the evaluation of synthetic varieties of maize for yield. *Crop Sci.* 8:448–451.
- Troyer, A. F., and A. R. Hallauer. 1968. Analysis of a diallel set of early flint varieties of maize. *Crop Sci.* 1:581–584.
- Hallauer, A. R. 1968. Estimates of genetic variances in Iowa Long Ear Synthetic, *Zea mays* L. *Advancing Frontiers Plant Sci.* 22:147–162.
- Hallauer, A. R., and J. H. Sears. 1969. Mass selection for yield in two varieties of maize. *Crop Sci.* 9:47–50.
- Hallauer, A. R., and J. H. Sears. 1969. Effect of thinning on stand, yield, and plant height in maize. *Crop Sci.* 9:514–515.
- Hallauer, A. R. 1970. Zygote selection for the development of single-cross hybrids in maize. *Advancing Frontiers Plant Sci.* 25:75–81.
- Klun, J. A., W. D. Guthrie, A. R. Hallauer, and W. A. Russell. 1970. The genetic nature of 2,4-dihydroxy-7-methoxy-1 4(2H) benzoxazin-3-one concentration and resistance to the European corn borer in a diallel set of eleven maize inbreds. *Crop Sci.* 10:87–90.
- Hallauer, A. R., and S. A. Eberhart. 1970. Reciprocal full-sib selection. *Crop Sci.* 10:315–316.

- Marquez-Sanchez, F., and A. R. Hallauer. 1970. Influence of sample size on the estimation of genetic variances in a synthetic variety of maize. I. Grain yield. *Crop Sci.* 10:357-361.
- Marquez-Sanchez, F., and A. R. Hallauer. 1970. Influence of sample size on the estimation of genetic variances in a synthetic variety of maize. II. Plant and ear characters. *Iowa State J. Sci.* 44:423-436.
- Hallauer, A. R. 1970. Genetic variability for yield after four cycles of reciprocal recurrent selection in maize. *Crop Sci.* 10:482-485.
- Hallauer, A. R. 1971. Change in genetic variance for seven plant and ear traits after four cycles of reciprocal recurrent selection for yield in maize. *Iowa State J. Sci.* 45:575-593.
- Burton, J. W., L. H. Penny, A. R. Hallauer, and S. A. Eberhart. 1971. Evaluation of synthetic populations developed from a maize variety (BSK) by two methods of recurrent selection. *Crop Sci.* 11:361-365.
- Wright, J. A., A. R. Hallauer, L. H. Penny, and S. A. Eberhart. 1971. Estimating genetic variance in maize by use of single and three-way crosses among unselected inbred lines. *Crop Sci.* 11:690-695.
- Hallauer, A. R. 1972. Third phase in the yield evaluation of synthetic varieties of maize. *Crop Sci.* 12:16-18.
- Eberhart, S. A., A. R. Hallauer, and W. A. Russell. 1972. Registration of a four maize germplasm synthetics. *Crop Sci.* 12:132.
- Hallauer, A. R., and J. H. Sears. 1972. Integrating exotic germplasm into Corn Belt maize breeding programs. *Crop Sci.* 12:203-206.
- Darrah, L. L., and A. R. Hallauer. 1972. Genetic effects estimated from generation means in four diallel sets of maize inbreds. *Crop Sci.* 12:615-621.
- Hallauer, A. R. 1972. Hybrid development and population improvement by full-sib selection. *Annu. Illinois Corn Breeders School.* 8:1-34.
- Hallauer, A. R., and A. F. Troyer. 1972. Prolific corn hybrids and minimizing risks of stress. *Annu. Hybrid Corn & Sorghum Res. Conf. Proc.* 26:140-158.
- Hallauer, A. R. 1973. Hybrid development and population improvement in maize by full-sib selection. *Egypt. J. Genet. Cytol.* 2:84-101.
- Hallauer, A. R., and J. H. Sears. 1973. Changes in quantitative traits associated with inbreeding in a synthetic variety of maize. *Crop Sci.* 13:327-330.
- Eberhart, S. A., Seme Debela, and A. R. Hallauer. 1973. Reciprocal recurrent selection in the BSSS and BSCB1 maize population and half-sib selection in BSSS. *Crop Sci.* 13:451-456.
- Hallauer, A. R. 1974. Heritability of prolificacy in maize. *J. Hered.* 65:163-168.
- Obilana, A. T., and A. R. Hallauer. 1974. Estimation of variability of quantitative traits in BSSS by using unselected maize inbred lines. *Crop Sci.* 14:99-103.
- Owens, J. C., D. C. Peters, and A. R. Hallauer. 1974. Corn rootworm tolerance in maize. *Environ. Entom.* 3:767-772.
- Hallauer, A. R., and C. A. Martinson. 1975. Reaction of maize hybrids and inbreds to *Bipolaris maydis*, Race T, infection. *Agron. J.* 67:497-501.
- Silva, J. C., and A. R. Hallauer. 1975. Estimation of epistatic variance in Iowa Stiff Stalk Synthetic variety of maize (*Zea mays* L.). *J. Hered.* 66:290-296.
- Hallauer, A. R., and C. A. Martinson. 1975. Material effects detected in maize hybrids infected with *Bipolaris maydis*. Race T. *Crop Sci.* 15:686-689.
- Hallauer, A. R. 1975. Relation of gene action and type of tester in maize breeding procedures. *Annu. Corn and Sorghum Res. Conf. Proc.* 29:150-165.
- Hallauer, A. R., and D. Malithano. 1976. Evaluation of maize varieties for their potential as breeding populations. *Euphytica* 25:117-127.
- Hoegemeyer, T. C., and A. R. Hallauer. 1976. Selection among and within full-sib families to develop single-crosses of maize. *Crop Sci.* 16:76-81.

- Martin, J. M., and A. R. Hallauer. 1976. Relation between heterozygosis and yield for four types of maize inbred lines. *Egypt. J. Genet. Cytol.* 5:119-135.
- Bartual, R., and A. R. Hallauer. 1976. Estimation of genetic variances in BSSS by use of inbred lines developed by full-sib selection. *Maydica* 21:49-60.
- Gama, Elto E. G., and A. R. Hallauer. 1977. Relation between inbred and hybrid traits in maize. *Crop Sci.* 17:703-706.
- Good, R. L., and A. R. Hallauer. 1977. Inbreeding depression in maize by selfing and full-sibbing. *Crop Sci.* 17:935-940.
- Obilana, A. T., and A. R. Hallauer. 1977. Comparison of estimates of genetic variability in Iowa Stiff Stalk Synthetic. *Maydica* 22:159-172.
- Hallauer, A. R., and G. F. Sprague. 1977. W. A. Russell (Dedication). *Maydica* 22:155-158.
- Guthrie, W. D., W. A. Russell, G. L. Reed, A. R. Hallauer, and D. F. Cox. 1978. Methods of evaluating maize for sheath-collar-feeding resistance to the European corn borer. *Maydica* 23:45-53.
- Russell, W. K., W. A. Russell, W. D. Guthrie, A. R. Hallauer, and J. C. Robins. 1978. Allocation of resources in breeding for resistance in maize to second brood of the European corn borer. *Maydica* 23:11-20.
- Miranda, J. B. Filho, and A. R. Hallauer. 1978. Correlation between population means in different generations under selfing and full-sibbing. One population. Brazil. *J. Genet.* 1:121-144.
- Hallauer, A. R. 1978. Introduction to cooperative state-federal corn breeding project at Ames, Iowa. *Annu. Illinois Corn Breeders School* 14:1-12.
- Hallauer, A. R. 1978. Recurrent selection programs. *Annu. Illinois Corn Breeders School.* 14:28-45.
- Cortez-Mendoza, H., and A. R. Hallauer. 1979. Divergent mass selection for ear length of maize. *Crop Sci.* 19:175-178.
- Obilana, A. T., A. R. Hallauer, and O. S. Smith. 1979. Estimated genetic variability in a maize interpopulation. *J. Hered.* 70:127-132.
- Obilana, A. T., A. R. Hallauer, and O. S. Smith. 1979. Predicted and observed response to reciprocal full-sib selection in maize (*Zea mays* L.). *Egypt. J. Genet. Cytol.* 8:269-282.
- Hallauer, A. R., and E. Lopez-Perez. 1979. Comparisons among testers for evaluating lines of corn. *Annu. Corn Sorghum Res. Conf. Proc.* 33:57-75.
- Martin, J. M., and A. R. Hallauer. 1980. Response from reciprocal recurrent selection in maize (*Zea mays* L.). *Crop Sci.* 20:599-603.
- Gama, Elto E. G., and A. R. Hallauer. 1980. Stability of hybrids produced from selected and unselected lines of maize. *Crop Sci.* 20:623-626.
- Hallauer, A. R. 1980. Relation of quantitative genetics and applied maize breeding. Brazil. *J. Genet.* 3:1-26.
- Robertson, D. S., A. R. Hallauer, and T. C. Kevern. 1981. Effects of substituting segments of the long arm of chromosome 10 from four inbred lines in the W22 inbred line of maize. *J. Hered.* 72:399-402.
- Lantin, M. M., and A. R. Hallauer. 1981. Response to reciprocal full-sib and mass selection in corn (*Zea mays* L.). *Iowa Acad. Sci.* 88:172-178.
- Smith, O. S., A. R. Hallauer, and W. A. Russell. 1981. Use of index selection in recurrent selection programs in maize. *Euphytica* 30:611-618.
- Smith, O. S., A. R. Hallauer, W. A. Russell, and T. M. Crosbie. 1981. Use of selection indices in maize improvement and hybrid development programs. *Proc. Annu. Corn Sorghum Res. Conf.* 36:95-103.

- Sadehdel-Moghaddan, M., P. J. Loesch, Jr., W. J. Wiser, and A. R. Hallauer. 1982. Interrelationships and inheritance of protein quality and agronomic traits in an opaque-2 synthetic of maize (*Zea mays* L.). *Iowa State J. Res.* 57:85–96.
- Moreno-Gonzalez, M., and A. R. Hallauer. 1982. Combined  $S_2$  and crossbred family selection in full-sib reciprocal recurrent selection. *Theor. Appl. Genet.* 61:355–358.
- Miranda, J. B. Filho, R. Vencovsky, and A. R. Hallauer. 1982. Correlation between population means under selfing and full-sibbing. Two populations. *Brazil. J. Genet.* 5:299–311.
- Kevern, T. C., and A. R. Hallauer. 1983. Relation of vertical root-pull resistance and maturity in maize populations. *Crop Sci.* 23:357–363.
- Sadehdel-Moghaddan, M., P. J. Loesch, Jr., A. R. Hallauer, and W. D. Guthrie. 1983. Inheritance of resistance to the first and second broods of the European corn borer in corn. *Iowa Acad. Sci.* 90:35–38.
- Mulamba, N. N., and A. R. Hallauer. 1983. Mass selection for grain yield and effects of three recurrent selection methods on genetic variability in a maize (*Zea mays* L.) population. *Crop Sci.* 13:536–540.
- Hallauer, A. R., W. A. Russell, and O. S. Smith. 1983. Quantitative analysis of Iowa Stiff Stalk Synthetic. *Proc. Stadler Genet. Sym.* 15:83–104.
- Silva, J., and A. R. Hallauer. 1984. Factors affecting production of corn forage. *Iowa Acad. Sci.* 91:76–81.
- Hallauer, A. R. 1984. Reciprocal full-sib selection in maize. *Crop Sci.* 24:755–759.
- Pego, S., and A. R. Hallauer. 1984. Performance of Portuguese maize germplasm with abnormal ear shape. *Maydica* 29:39–53.
- Hallauer, A. R. 1984. George F. Sprague: 60 years of corn research. *Maydica* 29:351–355.
- Lamkey, K. R., and A. R. Hallauer. 1984. Comparisons of populations improved by recurrent selection. *Maydica* 29:357–374.
- Oyervides-Garcia, M., A. R. Hallauer, and H. Cortez-Mendoza. 1985. Evaluation of improved corn populations in Mexico and the U.S. Corn Belt. *Crop Sci.* 25:115–120.
- Funnemark, W. L., and A. R. Hallauer. 1985. Cold tolerance in corn hybrids as affected by tillage systems and planting dates. *Iowa Acad. Sci.* 92:58–62.
- Hallauer, A. R., and T. S. Colvin. 1985. Response of corn hybrids to four methods of tillage. *Agron. J.* 77:547–550.
- Hallauer, A. R. 1985. Selection in mutation breeding. *Proc. Consultants Meeting—A review. Theor. Appl. Genet.* 64:76.
- Hallauer, A. R. 1985. Interaction of corn hybrids with tillage systems. *Illinois Corn Breeders School* 21:65–90.
- Gutierrez-Gaitan, M. A., H. Cortez-Mendoza, E. N. Wathika, C. O. Gardner, M. Oyervides-Garcia, A. R. Hallauer, and L. L. Darrah. 1986. Testcross evaluation of Mexican maize populations. *Crop Sci.* 26:99–104.
- Oyervides-Garcia, M., and A. R. Hallauer. 1986. Selection-included differences among strains of Iowa Stiff Stalk Synthetic maize. *Crop Sci.* 26:506–511.
- Salazar, A. M., and A. R. Hallauer. 1986. Divergent mass selection for ear length in maize. *Brazil. J. Genet.* 9:1–14.
- Clucas, C. P., and A. R. Hallauer. 1986. Effects of visual selection among and within  $S_1$  lines of maize on  $S_2$  line and testcross performance. *Iowa Acad. Sci.* 93:178–183.
- St. Martin, S. K., P. J. Loesch, Jr., and A. R. Hallauer. 1986. Heritabilities, correlations, and index selection in BSBB02 maize synthetic. *Maydica* 31:325–333.
- Lamkey, K. R., and A. R. Hallauer. 1986. Performance of high  $\times$  high, high  $\times$  low, and low  $\times$  low crosses of lines from the BSSS maize synthetic. *Crop Sci.* 26:1114–1118.

- Alexander, W. L., and A. R. Hallauer. 1986. Effect of one generation of self-pollination to increase seed supplies on predicted gain of some common recurrent selection schemes. *Brazil. J. Genet.* 9:545-548.
- Kahler, A. R., A. R. Hallauer, and C. O. Gardner. 1986. Allozyme polymorphisms within and among open-pollinated and adapted exotic populations of maize. *Theor. Appl. Genet.* 72:592-601.
- Price, S. C., A. L. Kahler, A. R. Hallauer, P. Charmley, and D. A. Giegel. 1986. Relationships between performance and multilocus heterozygosity at enzyme loci in single-cross hybrids of maize. *J. Hered.* 77:341-344.
- Lamkey, K. R., and A. R. Hallauer. 1987. Heritability estimates from recurrent selection experiments in maize. *Maydica* 32:61-78.
- Lamkey, K. R., A. R. Hallauer, and A. L. Kahler. 1987. Allelic differences at enzyme loci and hybrid performance in maize. *J. Hered.* 78:231-234.
- Reeder, L. R., Jr., A. R. Hallauer, and K. R. Lamkey. 1987. Estimation of genetic variability in two maize populations. *J. Hered.* 78:372-376.
- Rodriguez, O. A., and A. R. Hallauer. 1988. Effects of recurrent selection in corn populations. *Crop Sci.* 28:276-280.
- Lamkey, K. R., A. R. Hallauer, and D. S. Robertson. 1988. Contribution of the long arm of chromosome 10 to the total heterosis observed in five maize hybrids. *Crop Sci.* 28:896-901.
- Han, G. C., and A. R. Hallauer. 1988. Estimates of genet variability in  $F_2$  maize populations. *Iowa Acad. Sci.* 96:12-19.
- Kim, S.-K., J. L. Brewbaker, and A. R. Hallauer. 1988. Insect and disease resistance from tropical maize for use in temperate zone hybrids. *Proc. Corn Sorghum Ind. Res. Conf.* 43:194-226.
- Helms, T. C., A. R. Hallauer, and O. S. Smith. 1989. Genetic drift and selection evaluation from recurrent selection programs in maize. *Crop Sci.* 29:602-607.
- Kim, S.-K., A. R. Hallauer, W. D. Guthrie, D. Barry, K. R. Lamkey, and C. S. Hong. 1989. Genetic resistance of tropical inbreds to second generation European corn borer, *Ostrinia nubilalis*. *J. Econ. Entom.* 82:1207-1211.
- Kim, S.-K., W. D. Guthrie, A. R. Hallauer, W. A. Russell, J. L. Brewbaker, and C. S. Hong. 1989. Evaluation of tropical and subtropical corn lines for resistance to second-generation European corn borer. *J. Econ. Entom.* 82:1245-1250.
- Han, G.-C., A. R. Hallauer, and T. B. Bailey, Jr. 1989. Nonlinear relation between single-cross hybrids and their parental lines. *Brazil. J. Genet.* 12:287-301.
- Helms, T. C., A. R. Hallauer, and O. S. Smith. 1989. Genetic variability in improved and unimproved 'Iowa Stiff Stalk Synthetic' corn populations. *Crop Sci.* 29:959-962.
- Iglesias, C. A., and A. R. Hallauer. 1989. S2 recurrent selection in maize populations with exotic germplasm. *Maydica* 34:133-140.
- Kim, S. K., and A. R. Hallauer. 1989. Agronomic traits of tropical and subtropical maize inbreds in Iowa. *Plant Var. Seeds* 2:85-91.
- Covarrubias-Prieto, J., A. R. Hallauer, and K. R. Lamkey. 1989. Effects of intermating on means and variances of  $F_2$  populations of maize. *Genetika* 21:111-125.
- Hallauer, A. R. 1989. Improvement in yield of maize hybrids. *Svezak* 35:193-199.
- Ron-Parra, J., and A. R. Hallauer. 1989. Analyses of maize variety-environment variation in the Bajio region of Mexico. *Agricultura Tecnica Mexico* 35:3-22.
- Hallauer, A. R. 1989. Fifty years of recurrent selection in corn. *Illinois Corn Breeders School* 25:39-63.
- Hallauer, A. R. 1990. Methods used in developing maize lines. *Maydica* 36:1-16.



- Melchinger, A. E., M. Lee, K. R. Lamkey, A. R. Hallauer, and W. L. Woodman. 1990. Genetic diversity for restriction fragment length polymorphisms and heterosis for two diallel sets of maize inbreds. *Theor. Appl. Genet.* 80:488–496.
- Hallauer, A. R. 1990. Germplasm sources and breeding strategies for line development in the 1990's. *Proc. Corn Sorghum Res. Conf.* 45:64–79.
- Russell, J. R., N. A. Ilbreck, A. R. Hallauer, and D. R. Buxton. 1990. Nutritive value and ensiling characteristics of maize herbage as influenced by agronomic factors. *Animal Feed Sci. Tech.* 38:11–24.
- Lamkey, K. R., P. A. Peterson, and A. R. Hallauer. 1991. Frequency of the transposable element *Uq* in Iowa Stiff Stalk maize populations. *Genetical Res. Camb.* 57:1–9.
- Iglesias, C. A., and A. R. Hallauer. 1991. Response to S2 recurrent selection in exotic and semi-exotic populations of maize. *J. Iowa Acad. Sci.* 98:4–13.
- Rodriguez, O. A., and A. R. Hallauer. 1991. Variation among full-sib families of corn in different generations of inbreeding. *Crop Sci.* 31:43–47.
- Eyherabide, G. H., and A. R. Hallauer. 1991. Reciprocal full-sib selection in maize. I. Direct and indirect responses. *Crop Sci.* 31:952–959.
- Eyherabide, G. H., and A. R. Hallauer. 1991. Reciprocal full-sib selection in maize. II. Contributions of additive and dominance effects and effects of genetic drift. *Crop Sci.* 31:1442–1448.
- Valdivia-Bernal, R., and A. R. Hallauer. 1991. Estimates of genetic homeostasis in maize. *Brazil. J. Genet.* 14:483–499.
- de Rissi, R., and A. R. Hallauer. 1991. Evaluation of testers in a hybrid maize development program. *Brazil. J. Genet.* 14:467–481.
- Getschman, R. J., and A. R. Hallauer. 1991. Genetic variation among and within S<sub>1</sub> progenies of maize. *J. Iowa Acad. Sci.* 98:127–133.
- Stucker, D. S., and A. R. Hallauer. 1992. Genetic variability as affected by selection in Iowa Stiff Stalk Synthetic. *J. Hered.* 83:410–418.
- Graham, M. J., J. A. Hawk, R. B. Carroll, J. E. Ayers, K. R. Lamkey, and A. R. Hallauer. 1993. Evaluation of Iowa Stiff Stalk Synthetic for resistance to *Cercospora Zeae maydis*. *Plant Dis.* 77:382–385.
- Irlbeck, N. A., J. R. Russell, A. R. Hallauer, and D. R. Buxton. 1993. Nutritive value and ensiling characteristics of maize stover as influenced by hybrid maturity and generation, plant density and harvest date. *Animal Feed Sci. Tech.* 41:51–64.
- San Vicente, F. M., and A. R. Hallauer. 1993. Mass selection for adaptation in Antigua maize (*Zea mays* L.) composite. *J. Iowa Acad. Sci.* 100:9–12.
- San Vicente, F. M., and A. R. Hallauer. 1993. Inbreeding depression rates for two groups of maize inbred lines. *Brazil. J. Genet.* 16:989–1001.
- Benson, D. L., and A. R. Hallauer. 1994. Inbreeding depression rates in maize populations before and after recurrent selection. *J. Hered.* 85:122–128.
- Michellini, L. A., and A. R. Hallauer. 1993. Evaluation of exotic and adapted maize (*Zea mays* L.) germplasm crosses. *Maydica* 38:275–282.
- Lile, S. M., and A. R. Hallauer. 1994. Relation between S<sub>2</sub> and later generation testcrosses of two corn populations. *J. Iowa Acad. Sci.* 101:19–23.
- Lundvall, J. P., D. R. Buxton, A. R. Hallauer, and J. R. George. 1994. Forage quality variation among maize inbreds: in vitro digestibility and cell-wall components. *Crop Sci.* 34:1572–1578.
- Gama, Elto E. Gomes E, A. R. Hallauer, R. G. Ferro, and D. M. Barbosa. 1995. Heterosis in maize single crosses derived from a yellow Tuxpeno variety in Brazil. *Brazilian J. Genet.* 18:81–85.

- Gama, Elto E. Gomes E, A. R. Hallauer, M. A. Lopes, S. N. Parentoni, M. X. Dos Santos, and P. E. O. Guimaraes. 1995. Combining ability among fifteen early cycle maize populations in Brazil. *Brazilian J. Genet.* 18:569–577.
- Fountain, M. O., and A. R. Hallauer. 1996. Genetic variation within maize breeding populations. *Crop Sci.* 36:26–32.

## Books, Book Chapters, and Reviews

- Penny, L. H., W. A. Russell, G. F. Sprague, and A. R. Hallauer. 1963. Recurrent selection. p. 352–367. In: W. D. Hanson and H. F. Robinson (eds.), *Symp. Stat. Genet. Plant Breed.* NAS-NRC 982.
- Hallauer, A. R. 1978. Potential of exotic germplasm for maize improvement. p. 229–247. In: D. B. Walden (ed.), *Maize breeding and genetics*. Wiley, New York.
- Russell, W. A., and A. R. Hallauer. 1980. Corn. p. 299–312. In: W. R. Fehr and H. H. Hadley (eds.), *Hybridization of crop plants*. Am. Soc. Agron., Madison, WI.
- Hallauer, A. R. 1981. Selection and breeding methods. p. 1–35. In: K. J. Frey (ed.), *Plant breeding symp. II*. Iowa State Univ. Press, Ames.
- Hallauer, A. R., and J. B. Miranda Fo. 1981. Quantitative genetics in maize breeding. Iowa State Univ. Press, Ames (2nd edition, 1988).
- Hallauer, A. R. 1985. Compendium of recurrent selection methods and their application. *Critical Rev. Plant Sci.* 3:1–34.
- Hallauer, A. R. 1987. Selection methods used in the improvement of maize. In: A. S. Khehra, B. S. Brar, and D. S. Dhillon (eds.), *Plant breeding: today and tomorrow*. Ludhiani, India.
- Hallauer, A. R. 1987. Recurrent selection. p. 3–29. In: B. S. Ghai (ed.), *Plant genetics breeding review*. Ludhiari, India.
- Hallauer, A. R. 1987. Maize. p. 249–294. In: W. R. Fehr (ed.), *Principles of cultivar development*. Macmillan, New York.
- Hallauer, A. R. 1987. Breeding systems. p. 61–91. In: B. R. Christie (ed.), *Handbook of plant science in agriculture*. CRC Press, Boca Raton, FL.
- Hallauer, A. R., W. A. Russell, and K. R. Lamkey. 1988. Corn breeding. p. 463–564. In: G. F. Sprague and J. W. Dudley (eds.), *Corn and corn improvement*. Am. Soc. Agron., Madison, WI.
- Hallauer, A. R. 1992. Recurrent selection in maize. *Plant Breed. Rev.* 9: 115–179.
- Hallauer, A. R. 1994. Corn genetics and breeding. *Encyclopedia of Agricultural Science* 1:455–467. Academic Press, New York.
- Hallauer, A. R. (ed.) 1994. *Specialty corns*. CRC Press, Boca Raton, FL.
- Hallauer, A. R. 1995. International activities in maize germplasm. p. 149–163. In: R. R. Duncan (ed.), *International germplasm transfer: past and present*. CSSA Spec. Publ. 23. ASA, CSSA, and SSSA, Madison, WI.
- Ron-Parra, J., and A. R. Hallauer. 1996. Utilization of tropical-subtropical maize germplasm. *Plant Breed. Rev.* 14:165–187.

## Germplasm Released

- Russell, W. A., L. H. Penny, A. R. Hallauer, G. E. Scott, W. D. Guthrie, and F. F. Dicke. 1971. Registration of maize (GP 6 to 21) germplasm synthetics. *Crop Sci.* 11:140–141.
- Hallauer, A. R., S. A. Eberhart, and W. A. Russell. 1974. Registration of maize germplasm (No. GP 26 to 34). *Crop Sci.* 14:341–342.

- Russell, W. A., and A. R. Hallauer. 1974. Registration of B76 and B77 parental lines of maize. *Crop Sci.* 14:778.
- Russell, W. A., and A. R. Hallauer. 1976. Registration of B79 parental line of maize (*Zea mays* L.) (Reg. No. PL 42). *Crop Sci.* 16:316.
- Hallauer, A. R., and O. S. Smith. 1979. Registration of maize germplasm (Reg. No. GP 81 and GP 82). *Crop Sci.* 19:755.
- Hallauer, A. R. 1986. Registration of BS26 germplasm. *Crop Sci.* 26:839.
- Hallauer, A. R., and W. A. Russell. 1986. Registration of BS18 germplasm. *Crop Sci.* 26:838.
- Hallauer, A. R. 1992. Registration of BS27 maize germplasm. *Crop Sci.* 32:1512–1513.
- Hallauer, A. R., K. R. Lamkey, W. A. Russell, and P. R. White. 1992. Registration of B95 parental inbred line of maize. *Crop Sci.* 32:1515.
- Hallauer, A. R. 1994. Registration of BS28 and BS29 maize germplasm. *Crop Sci.* 34:544–545.
- Hallauer, A. R., K. R. Lamkey, W. A. Russell, and P. R. White. 1994. Registration of B97 and B98 parental lines of maize. *Crop Sci.* 34: 318–319.
- Hallauer, A. R. 1995. Registration of BS30 maize germplasm. *Crop Sci.* 35:1234.
- Hallauer, A. R., and A. D. Wright. 1995. Registration of B101 germplasm line of maize. 35:1238–1239.
- Hallauer, A. R., K. R. Lamkey, W. A. Russell, and P. R. White. 1995. Registration of B99 and B100 inbred lines of maize. *Crop Sci.* 35:1714–1715.